

Resolving it into partial fraction:

Question # 1

$$\frac{1}{x^2 - 1}$$

Solution

$$\frac{1}{x^2 - 1} = \frac{1}{(x-1)(x+1)}$$

Now suppose

$$\frac{1}{(x-1)(x+1)} = \frac{A}{x-1} + \frac{B}{x+1}$$

Multiplying both sides by $(x-1)(x+1)$ we get

$$1 = A(x+1) + B(x-1) \dots \dots \dots \text{(i)}$$

Put $x-1=0 \Rightarrow x=1$ in equation (i)

$$1 = A(1+1) + B(0) \Rightarrow 1 = 2A + 0 \Rightarrow A = \frac{1}{2}$$

Now put $x+1=0 \Rightarrow x=-1$ in equation (i)

$$1 = A(0) + B(-1 - 1) \Rightarrow 1 = 0 - 2B \Rightarrow B = -\frac{1}{2}$$

Hence

$$\frac{1}{(x-1)(x+1)} = \frac{A}{x-1} + \frac{B}{x+1}$$

Question # 2

$$\frac{x^2(x^2 + 1)}{(x + 1)(x - 1)}$$

Solution

$$\begin{aligned} \frac{x^2(x^2+1)}{(x+1)(x-1)} &= \frac{x^4+x^2}{(x^2-1)} \\ &= x^2 + 2 + \frac{2}{(x^2-1)} = x^2 + 2 + \frac{2}{(x+1)(x-1)} \end{aligned}$$

Now consider

$$\frac{2}{(x+1)(x-1)} = \frac{A}{x+1} + \frac{B}{x-1}$$

Multiplying both sides by $(x+1)(x-1)$

$$2 = A(x-1) + B(x+1) \quad \dots \dots \dots \text{(i)}$$

Put $x + 1 = 0 \Rightarrow x = -1$ in equation (i)

$$2 = A(-1-1) + B(0) \Rightarrow 2 = -2A + 0 \Rightarrow A = -1$$

Now put $x-1=0 \Rightarrow x=1$ in equation (i)

$$2 = A(0) + B(1+1) \Rightarrow 2 = 0 + 2B \Rightarrow B = 1$$

$$\text{So} \quad \frac{2}{(x+1)(x-1)} = \frac{-1}{x+1} + \frac{1}{x-1}$$

Hence

$$\begin{aligned} \frac{x^2(x^2+1)}{(x+1)(x-1)} &= x^2 + 2 + \frac{-1}{(x+1)} + \frac{1}{(x-1)} \\ &= x^2 + 2 - \frac{1}{(x+1)} + \frac{1}{(x-1)} \end{aligned} \quad Answer$$

Question # 3

$$\frac{2x+1}{(x-1)(x+2)(x+3)}$$

Solution

$$\frac{2x+1}{(x-1)(x+2)(x+3)} = \frac{A}{x-1} + \frac{B}{x+2} + \frac{C}{x+3}$$

Multiplying both side by $(x-1)(x+2)(x+3)$

Put $x-1=0 \Rightarrow x=1$ in equation (i)

$$2(1) + 1 = A(1+2)(1+3) + B(0) + C(0)$$

$$3 = A(3)(4) + 0 + 0 \quad \Rightarrow \quad 3 = 12A \quad \Rightarrow \quad \frac{3}{12} = A \quad \Rightarrow \quad \boxed{A = \frac{1}{4}}$$

Now put $x + 2 = 0 \Rightarrow x = -2$ in equation (i)

$$2(-2) + 1 = A(0) + B(-2 - 1)(-2 + 3) + C(0)$$

$$-4 + 1 = 0 + B(-3)(1) + 0 \quad \Rightarrow \quad -3 = -3B \quad \Rightarrow \quad B = 1$$

Now put $x + 3 = 0 \Rightarrow x = -3$ in equation (i)

$$2(-3) + 1 = A(0) + B(0) + C(-3 - 1)(-3 + 2)$$

$$-6 + 1 = 0 + 0 + C(-4)(-1) \quad \Rightarrow \quad -5 = 4C \quad \Rightarrow \quad C = -\frac{5}{4}$$

So

$$\begin{aligned} \frac{2x+1}{(x-1)(x+2)(x+3)} &= \frac{1/4}{x-1} + \frac{1}{x+2} + \frac{-5/4}{x+3} \\ &= \frac{1}{4(x-1)} + \frac{1}{x+2} - \frac{5}{4(x+3)} \end{aligned} \quad \text{Answer}$$

Question # 4

$$\frac{3x^2 - 4x - 5}{(x-2)(x^2 + 7x + 10)} \quad \because x^2 + 7x + 10 = x^2 + 5x + 2x + 10 \\ = x(x+5) + 2(x+5)$$

Solution

$$\frac{3x^2 - 4x - 5}{(x-2)(x^2 + 7x + 10)} = \frac{3x^2 - 4x - 5}{(x-2)(x+5)(x+2)}$$

Now resolving into partial fraction.

$$\frac{3x^2 - 4x - 5}{(x-2)(x+5)(x+2)} = \frac{A}{x-2} + \frac{B}{x+5} + \frac{C}{x+2}$$

$$\left[\begin{array}{l} \text{Do yourself. You will get} \\ A = -\frac{1}{28}, B = \frac{30}{7}, C = -\frac{5}{4} \end{array} \right]$$

Question # 5

$$\frac{1}{(x-1)(2x-1)(3x-1)}$$

Solution

$$\frac{1}{(x-1)(2x-1)(3x-1)} = \frac{A}{x-1} + \frac{B}{2x-1} + \frac{C}{3x-1}$$

Multiplying both side by $(x-1)(2x-1)(3x-1)$.

$$1 = A(2x-1)(3x-1) + B(x-1)(3x-1) + C(2x-1)(3x-1) \dots \dots \dots \text{(i)}$$

Put $x-1=0 \Rightarrow x=1$ in equation (i)

$$1 = A(2(1)-1)(3(1)-1) + B(0) + C(0) \Rightarrow 1 = A(1)(2) + 0 + 0$$

$$\Rightarrow 1 = 2A \Rightarrow \boxed{A = \frac{1}{2}}$$

Put $2x-1=0 \Rightarrow 2x=1 \Rightarrow x=\frac{1}{2}$ in equation (i)

$$1 = A(0) + B\left(\frac{1}{2}-1\right)\left(3\left(\frac{1}{2}\right)-1\right) + C(0) \Rightarrow 1 = 0 + B\left(-\frac{1}{2}\right)\left(\frac{1}{2}\right) + 0$$

$$\Rightarrow 1 = -\frac{1}{4}B \Rightarrow \boxed{B = -4}$$

Put $3x-1=0 \Rightarrow 3x=1 \Rightarrow x=\frac{1}{3}$ in equation (i)

$$1 = A(0) + B(0) + C\left(\frac{1}{3}-1\right)\left(2\left(\frac{1}{3}\right)-1\right) \Rightarrow 1 = 0 + 0 + C\left(-\frac{2}{3}\right)\left(-\frac{1}{3}\right)$$

$$\Rightarrow 1 = \frac{2}{9}C \Rightarrow \boxed{C = \frac{9}{2}}$$

Hence

$$\begin{aligned}\frac{1}{(x-1)(2x-1)(3x-1)} &= \frac{\cancel{1}/2}{x-1} + \frac{-4}{2x-1} + \frac{\cancel{9}/2}{3x-1} \\ &= \frac{1}{2(x-1)} - \frac{4}{2x-1} + \frac{9}{2(3x-1)} \quad \text{Answer}\end{aligned}$$

Question # 6

$$\frac{x}{(x-a)(x-b)(x-c)}$$

Solution

$$\frac{x}{(x-a)(x-b)(x-c)} = \frac{A}{x-a} + \frac{B}{x-b} + \frac{C}{x-c}$$

Multiplying both sides by $(x-a)(x-b)(x-c)$.

$$x = A(x-b)(x-c) + B(x-a)(x-c) + C(x-a)(x-b) \dots \dots \dots \text{(i)}$$

Put $x-a=0 \Rightarrow x=a$ in equation (i)

$$a = A(a-b)(a-c) + B(0) + C(0)$$

$$\Rightarrow a = A(a-b)(a-c) + 0 + 0 \Rightarrow A = \frac{a}{(a-b)(a-c)}$$

Now put $x-b=0 \Rightarrow x=b$ in equation (i)

$$a = A(0) + B(b-a)(b-c) + C(0)$$

$$\Rightarrow a = 0 + B(b-a)(b-c) + 0 \Rightarrow B = \frac{b}{(b-a)(b-c)} \quad \text{Now put}$$

$x-c=0 \Rightarrow x=c$ in equation (i)

$$c = A(0) + B(0) + C(c-a)(c-b)$$

$$\Rightarrow c = 0 + 0 + C(c-a)(c-b) \Rightarrow C = \frac{c}{(c-a)(c-b)}$$

So

$$\begin{aligned}\frac{x}{(x-a)(x-b)(x-c)} &= \frac{\cancel{a}/(a-b)(a-c)}{x-a} + \frac{\cancel{b}/(b-a)(b-c)}{x-b} + \frac{\cancel{c}/(c-a)(c-b)}{x-c} \\ &= \frac{a}{(a-b)(a-c)(x-a)} + \frac{b}{(b-a)(b-c)(x-b)} + \frac{c}{(c-a)(c-b)(x-c)}\end{aligned}$$

Answer

Question # 7

$$\frac{6x^3 + 5x^2 - 7}{2x^2 - x - 1}$$

Solution

$$\frac{6x^3 + 5x^2 - 7}{2x^2 - x - 1}$$

$$\begin{aligned}
 &= 3x+4 + \frac{7x-3}{2x^2-x-1} \\
 &= 3x+4 + \frac{7x-3}{2x^2-2x+x-1} \\
 &= 3x+4 + \frac{7x-3}{2x(x-1)+1(x-1)} \\
 &= 3x+4 + \frac{7x-3}{(x-1)(2x+1)}
 \end{aligned}$$

$$\begin{array}{r}
 & 3x+4 \\
 2x^2 - x - 1 & \overline{)6x^3 + 5x^2 - 7} \\
 & - 6x^3 - 3x^2 - 3x \\
 & + + \\
 & 8x^2 + 3x - 7 \\
 & - 8x^2 - 4x - 4 \\
 & + + \\
 & 7x - 3
 \end{array}$$

Now Consider

$$\frac{7x-3}{(x-1)(2x+1)} = \frac{A}{x-1} + \frac{B}{2x+1}$$

Find value of A & B yourself
 You will get A = $\frac{4}{3}$ and B = $\frac{13}{3}$

so

$$\frac{7x-3}{(x-1)(2x+1)} = \frac{\frac{4}{3}}{x-1} + \frac{\frac{13}{3}}{2x+1} = \frac{4}{3(x-1)} + \frac{13}{3(2x+1)}$$

Hence

$$\frac{6x^3+5x^2-7}{2x^2-x-1} = 3x+4 + \frac{4}{3(x-1)} + \frac{13}{3(2x+1)} \quad \text{Answer}$$

Question # 8

$$\begin{array}{c}
 \frac{2x^3+x^2-5x+3}{2x^3+x^2-3x} \\
 \frac{2x^3+x^2-5x+3}{2x^3+x^2-3x} \\
 = 1 + \frac{-2x+3}{2x^3+x^2-3x} \\
 = 1 + \frac{-2x+3}{x(2x^2+x-3)} \\
 = 1 + \frac{-2x+3}{x(x(2x+3)-1(2x+3))} \\
 \frac{1}{2x^3+x^2-3x} \\
 \frac{2x^3+x^2-3x}{-2x+3}
 \end{array}$$

Now consider

$$\frac{3-2x}{x(2x+3)(x-1)} = \frac{A}{x} + \frac{B}{2x+3} + \frac{C}{x-1}$$

$$\Rightarrow 3-2x = A(2x+3)(x-1) + Bx(x-1) + Cx(2x+3) \dots \dots \dots \text{(i)}$$

Put $x=0$ in equation (i)

$$3-2(0) = A(2(0)+3)((0)-1) + B(0) + C(0) \Rightarrow 3-0 = A(0+3)(-1) + 0 + 0$$

$$\Rightarrow 3 = -3A \Rightarrow A = -1$$

Now put $2x+3=0 \Rightarrow 2x=-3 \Rightarrow x=-\frac{3}{2}$ in equation (i)

$$3 - 2\left(-\frac{3}{2}\right) = A(0) + B\left(-\frac{3}{2}\right)\left(-\frac{3}{2} - 1\right) + C(0) \Rightarrow 3 + 3 = 0 + B\left(-\frac{3}{2}\right)\left(-\frac{5}{2}\right) + 0$$

$$\Rightarrow 6 = \frac{15}{4}B \Rightarrow B = (6)\left(\frac{4}{15}\right) \Rightarrow B = \frac{8}{5}$$

Now put $x-1=0 \Rightarrow x=1$ in equation (i)

$$3 - 2(1) = A(0) + B(0) + C(1)(2(1) + 3) \Rightarrow 1 = 0 + 0 + 5C \Rightarrow C = \frac{1}{5}$$

So $\frac{3-2x}{x(2x+3)(x-1)} = \frac{-1}{x} + \frac{\frac{8}{5}}{2x+3} + \frac{\frac{1}{5}}{x-1} = -\frac{1}{x} + \frac{8}{5(2x+3)} + \frac{1}{5(x-1)}$

Hence $\frac{2x^3+x^2-5x+3}{2x^3+x^2-3x} = 1 - \frac{1}{x} + \frac{8}{5(2x+3)} + \frac{1}{5(x-1)}$ Answer

Question # 9

$$\frac{(x-1)(x-3)(x-5)}{(x-2)(x-4)(x-6)}$$

Solution

$$\begin{aligned} \frac{(x-1)(x-3)(x-5)}{(x-2)(x-4)(x-6)} &= \frac{(x-1)(x^2-3x-5x+15)}{(x-2)(x^2-4x-6x+24)} \\ &= \frac{(x-1)(x^2-8x+15)}{(x-2)(x^2-10x+24)} = \frac{x^3-8x^2+15x-x^2+8x-15}{x^3-10x^2+24x-2x^2+20x-48} \\ &= \frac{x^3-9x^2+23x-15}{x^3-12x^2+44x-48} \\ &= 1 + \frac{3x^2-21x+33}{x^3-12x^2+44x-48} && x^3-12x^2+44x-48 \Big| x^3-9x^2+23x-15 \\ &= 1 + \frac{3x^2-21x+33}{(x-2)(x-4)(x-6)} && \frac{x^3-12x^2+44x-48}{-} \frac{x^3-9x^2+23x-15}{+} \\ &&& \hline &&& 3x^2-21x+33 \end{aligned}$$

Now Suppose

$$\frac{3x^2-21x+33}{(x-2)(x-4)(x-6)} = \frac{A}{x-2} + \frac{B}{x-4} + \frac{C}{x-6}$$

Find value of A, B and C yourself
You will get A = 3/8, B = 3/4, C = 15/8

So $\frac{3x^2-21x+33}{(x-2)(x-4)(x-6)} = \frac{\frac{3}{8}}{x-2} + \frac{\frac{3}{4}}{x-4} + \frac{\frac{15}{8}}{x-6}$

$$= \frac{3}{8(x-2)} + \frac{3}{4(x-4)} + \frac{15}{8(x-6)}$$

Hence

$$\frac{(x-1)(x-3)(x-5)}{(x-2)(x-4)(x-6)} = 1 + \frac{3}{8(x-2)} + \frac{3}{4(x-4)} + \frac{15}{8(x-6)} \quad Answer$$

Question # 10

$$\frac{1}{(1-ax)(1-bx)(1-cx)}$$

Solution

$$\frac{1}{(1-ax)(1-bx)(1-cx)} = \frac{A}{1-ax} + \frac{B}{1-bx} + \frac{C}{1-cx}$$

Multiplying both sides by $(1 - ax)(1 - bx)(1 - cx)$.

Put $1 - ax = 0 \Rightarrow ax = 1 \Rightarrow x = \frac{1}{a}$ in equation (i).

$$1 = A \left(1 - b \cdot \frac{1}{a} \right) \left(1 - c \cdot \frac{1}{a} \right) + B(0) + C(0) \quad \Rightarrow \quad 1 = A \left(1 - \frac{b}{a} \right) \left(1 - \frac{c}{a} \right) + 0 + 0$$

$$\Rightarrow 1 = A \left(\frac{a-b}{a} \right) \left(\frac{a-c}{a} \right) \quad \Rightarrow \quad 1 = A \frac{(a-b)(a-c)}{a^2} \quad \Rightarrow \quad A = \frac{a^2}{(a-b)(a-c)}$$

Find value of B & C yourself as A.

You will get $B = \frac{b^2}{(b-a)(b-c)}$, $C = \frac{c^2}{(c-a)(c-b)}$

$$\text{Hence } \frac{1}{(1-ax)(1-bx)(1-cx)} = \frac{\frac{a^2}{(a-b)(a-c)}}{1-ax} + \frac{\frac{b^2}{(b-a)(b-c)}}{1-bx} + \frac{\frac{c^2}{(c-a)(c-b)}}{1-cx}$$

$$= \frac{a^2}{(a-b)(a-c)(1-ax)} + \frac{b^2}{(b-a)(b-c)(1-bx)} + \frac{c^2}{(c-a)(c-b)(1-cx)}$$

Answer

Question # 11

$$\frac{x^2 + a^2}{(x^2 + b^2)(x^2 + c^2)(x^2 + d^2)}$$

$$\frac{x^2 + a^2}{(x^2 + b^2)(x^2 + c^2)(x^2 + d^2)}$$

Put $y = x^2$ in above.

$$\frac{y+a^2}{(y+b^2)(y+c^2)(y+d^2)}$$

Now consider

$$\frac{y+a^2}{(y+b^2)(y+c^2)(y+d^2)} = \frac{A}{y+b^2} + \frac{B}{y+c^2} + \frac{C}{y+d^2}$$

Put $y+b^2=0 \Rightarrow y=-b^2$ in equation (i)

$$-b^2+a^2=A(-b^2+c^2)(-b^2+d^2)+B(0)+C(0)$$

$$\Rightarrow a^2-b^2=A(c^2-b^2)(d^2-b^2)+0+0 \Rightarrow A=\frac{a^2-b^2}{(c^2-b^2)(d^2-b^2)}$$

Now put $y+c^2=0 \Rightarrow y=-c^2$ in equation (i)

$$-c^2+a^2=A(0)+B(-c^2+b^2)(-b^2+d^2)+C(0)$$

$$\Rightarrow a^2-c^2=0+B(b^2-c^2)(d^2-c^2)+0 \Rightarrow B=\frac{a^2-c^2}{(b^2-c^2)(d^2-c^2)}$$

Now put $y+d^2=0 \Rightarrow y=-d^2$ in equation (i)

$$-d^2+a^2=A(0)+B(0)+C(-d^2+b^2)(-d^2+c^2)$$

$$\Rightarrow a^2-d^2=0+0+C(b^2-d^2)(c^2-d^2) \Rightarrow C=\frac{a^2-d^2}{(b^2-d^2)(c^2-d^2)}$$

Hence

$$\begin{aligned} \frac{y+a^2}{(y+b^2)(y+c^2)(y+d^2)} &= \frac{\frac{a^2-b^2}{(c^2-b^2)(d^2-b^2)}}{y+b^2} + \frac{\frac{a^2-c^2}{(b^2-c^2)(d^2-c^2)}}{y+c^2} + \frac{\frac{a^2-d^2}{(b^2-d^2)(c^2-d^2)}}{y+d^2} \\ &= \frac{a^2-b^2}{(c^2-b^2)(d^2-b^2)(y+b^2)} + \frac{a^2-c^2}{(b^2-c^2)(d^2-c^2)(y+c^2)} + \frac{a^2-d^2}{(b^2-d^2)(c^2-d^2)(y+d^2)} \end{aligned}$$

Since $y=x^2$

$$= \frac{a^2-b^2}{(c^2-b^2)(d^2-b^2)(x^2+b^2)} + \frac{a^2-c^2}{(b^2-c^2)(d^2-c^2)(x^2+c^2)} + \frac{a^2-d^2}{(b^2-d^2)(c^2-d^2)(x^2+d^2)}$$

If you found any error, please report us at www.megalecture.com/pages/contact.html



Available online at <http://www.megalecture.com> in PDF Format (Picture format to view online). Page setup: A4 (8.27