

## MOLES AND STOICHIOMETRY WS 1

- 1 Nickel makes up 20% of the total mass of a coin. The coin has a mass of 10.0g.

How many nickel atoms are in the coin?


- A  $2.05 \times 10^{22}$     B  $4.30 \times 10^{22}$     C  $1.03 \times 10^{23}$     D  $1.20 \times 10^{24}$

$$\eta_{\text{nickel}} = \frac{2}{58.9} = 0.033 \text{ mol}$$

$$\# \text{ Atoms} = 0.033 \times 6.02 \times 10^{23} = 2.044 \times 10^{22}$$

- 2 On collision, airbags in cars inflate rapidly due to the production of nitrogen.

The nitrogen is formed according to the following equations.



How many moles of nitrogen gas are produced from 1 mol of sodium azide,  $\text{NaN}_3$ ?

- A 1.5    B 1.6    C 3.2    D 4.0

- 3 What is the number of molecules in  $500 \text{ cm}^3$  of oxygen under room conditions?

- A  $1.25 \times 10^{22}$   
 B  $1.34 \times 10^{22}$   
 C  $3.0 \times 10^{22}$   
 D  $3.0 \times 10^{26}$

$$\# \text{ O}_2 \text{ molecules} = \eta \times L$$

$$= \frac{500}{24000} \times 6.02 \times 10^{23}$$

$$= 1.25 \times 10^{22}$$

They're trying to confuse you.

- 4 Analytical chemists can detect very small amounts of amino acids, down to  $3 \times 10^{-21} \text{ mol}$ . How many molecules of an amino acid ( $M_r = 200$ ) would this be?

- A 9    B 200    C 1800    D 360 000

$$\text{Atoms} = \eta \times L$$

$$= 3 \times 10^{-21} \times 6.02 \times 10^{23} = 1806$$

- 5 Which of these samples of gas contains the same number of atoms as 1g of hydrogen ( $M_r : \text{H}_2, 2$ )?

- A 22g of carbon dioxide ( $M_r : \text{CO}_2, 44$ )  
 B 8g of methane ( $M_r : \text{CH}_4, 16$ )  
 C 20g of neon ( $M_r : \text{Ne}, 20$ )  
 D 8g of ozone ( $M_r : \text{O}_3, 48$ )

$$\eta_{\text{H}_2} = \frac{1\text{g}}{2} = 0.5 \text{ mol}$$

$$\eta_{\text{H atoms}} = 0.5 \times 2 = 1 \text{ mol}$$

$$\eta = \frac{20}{20} = 1 \text{ mol}$$

- 6 Which mass of gas would occupy a volume of  $3 \text{ dm}^3$  at  $25^\circ\text{C}$  and 1 atmosphere pressure? [1 mol of gas occupies  $24 \text{ dm}^3$  at  $25^\circ\text{C}$  and 1 atmosphere pressure.]

- A 3.2g  $\text{O}_2$  gas  
 B 5.6g  $\text{N}_2$  gas  
 C 8.0g  $\text{SO}_2$  gas  
 D 11.0g  $\text{CO}_2$  gas


$$3.2/32 = 0.1 \text{ mol}$$

$$5.6/28 = 0.2 \text{ mol}$$

$$8/64 = 0.125 \text{ mol}$$

$$\eta = \frac{3}{24} = 0.125 \text{ mol}$$

- 7 Most modern cars are fitted with airbags. These work by decomposing sodium azide to liberate nitrogen gas, which inflates the bag.



$$0.769 : 1.534 \text{ mol}$$

A typical driver's airbag contains 50 g of sodium azide.

Calculate the volume of nitrogen this will produce at room temperature.

- A 9.2 dm<sup>3</sup>      B 13.9 dm<sup>3</sup>      **C 27.7 dm<sup>3</sup>**      D 72.0 dm<sup>3</sup>

$$\downarrow \times 24 \text{ dm}^3 \text{ mol}^{-1}$$

$$27.69 \text{ dm}^3$$

- 8 N<sub>2</sub>O<sub>4</sub> is a poisonous gas. It can be disposed of safely by reaction with sodium hydroxide.  $n_{\text{NaOH}} = 2 \times 0.02 = 0.04 \text{ mol}$




What is the minimum volume of 0.5 mol dm<sup>-3</sup> NaOH(aq) needed to dispose of 0.02 mol of N<sub>2</sub>O<sub>4</sub>?

- A 8 cm<sup>3</sup>      B 12.5 cm<sup>3</sup>      C 40 cm<sup>3</sup>      **D 80 cm<sup>3</sup>**

$$V = \frac{n}{c} = \frac{0.04}{0.5} = 0.08$$

- 9 A household bleach contains sodium chlorate(I), NaClO, as its active ingredient. The concentration of NaClO in the bleach can be determined by reacting a known amount with aqueous hydrogen peroxide, H<sub>2</sub>O<sub>2</sub>.



When 25.0 cm<sup>3</sup> of bleach is treated with an excess of aqueous H<sub>2</sub>O<sub>2</sub>, 0.0350 mol of oxygen gas is given off.

What is the concentration of NaClO in the bleach?

$$c = \frac{0.035}{0.025} = 1.4 \text{ dm}^{-3}$$

- A  $8.75 \times 10^{-4} \text{ mol dm}^{-3}$   
 B 0.700 mol dm<sup>-3</sup>  
 C 0.875 mol dm<sup>-3</sup>  
**D 1.40 mol dm<sup>-3</sup>**



$$\text{M}_r : 151.7 : 79.9$$

$$19 : x$$

- 10 Titanium(IV) oxide, TiO<sub>2</sub>, is brilliantly white and much of the oxide produced is used in the manufacture of paint.

What is the maximum amount of TiO<sub>2</sub> obtainable from 19.0 tonnes of the ore ilmenite, FeTiO<sub>3</sub>?

- A 10.0 tonnes**      B 12.7 tonnes      C 14.0 tonnes      D 17.7 tonnes

$$x = 10$$

- 11 The foul smell that skunks spray is due to a number of thiols, one of which is methanethiol, CH<sub>3</sub>SH, which burns as follows.



A sample of 10 cm<sup>3</sup> of methanethiol was exploded with 60 cm<sup>3</sup> of oxygen.

What would be the final volume of the resultant mixture of gases when cooled to room temperature?




- A 20 cm<sup>3</sup>      B 30 cm<sup>3</sup>      C 50 cm<sup>3</sup>      D 70 cm<sup>3</sup>

12 The reaction between aluminium powder and anhydrous barium nitrate is used as the propellant in some fireworks. The metal oxides and nitrogen are the only products.

Which volume of nitrogen, measured under room conditions, is produced when 0.783 g of anhydrous barium nitrate reacts with an excess of aluminium?

- A 46.8 cm<sup>3</sup>    B 72.0 cm<sup>3</sup>    C 93.6 cm<sup>3</sup>    D 144 cm<sup>3</sup>

13 The amount of calcium ions in a sample of natural water can be determined by using an ion-exchange column as shown in the diagram.



③  $C = \frac{1.25 \times 10^{-4}}{0.05} = 2.5 \times 10^{-3} \text{ mol}$

A 50 cm<sup>3</sup> sample of water containing dissolved calcium sulphate was passed through the ion-exchange resin. Each calcium ion in the sample was exchanged for two hydrogen ions. The resulting acidic solution collected in the flask required 25 cm<sup>3</sup> of 1.0 × 10<sup>-2</sup> mol dm<sup>-3</sup> potassium hydroxide for complete neutralisation.

What was the concentration of the calcium sulphate in the original sample?

- A 2.5 × 10<sup>-3</sup> mol dm<sup>-3</sup>  
 B 1.0 × 10<sup>-2</sup> mol dm<sup>-3</sup>  
 C 2.0 × 10<sup>-2</sup> mol dm<sup>-3</sup>  
 D 4.0 × 10<sup>-2</sup> mol dm<sup>-3</sup>

$\frac{12 \times 8}{29 \times 4 + 207.2}$

14 Tetraethyl lead, Pb(C<sub>2</sub>H<sub>5</sub>)<sub>4</sub>, has been used as a petrol additive.

What is the percentage by mass of carbon in tetraethyl lead?

- A 10.2    B 14.9    C 29.7    D 32.0

- 15 A piece of rock has a mass of 2.00g. It contains calcium carbonate, but no other basic substances. It neutralises exactly 36.0 cm<sup>3</sup> of 0.500 mol dm<sup>-3</sup> hydrochloric acid.

What is the percentage of calcium carbonate in the 2.00 g piece of rock?

- A 22.5%      B 45.0%      C 72.0%      D 90.1%

- 16 In China, the concentration of blood glucose, C<sub>6</sub>H<sub>12</sub>O<sub>6</sub>, is measured in mmol/l. In Pakistan, the concentration of blood glucose is measured in mg/dl.

The unit l is a litre (1 dm<sup>3</sup>). The unit dl is a decilitre (0.1 dm<sup>3</sup>).

A blood glucose concentration of 18.5 mmol/l indicates a health problem. →

What is 18.5 mmol/l converted to mg/dl?

- A 33.3 mg/dl      B 178 mg/dl      C 333 mg/dl      D 3330 mg/dl

- 17 A 0.005 mol sample of anhydrous calcium carbonate was completely thermally decomposed to give 100 cm<sup>3</sup> of gas measured at a certain temperature and pressure.

In a separate experiment carried out at the same temperature and pressure, a 0.005 mol sample of anhydrous calcium nitrate was completely thermally decomposed. The volume of gaseous products was measured.

What total volume of gaseous products was produced from the calcium nitrate?

- A 50 cm<sup>3</sup>      B 100 cm<sup>3</sup>      C 200 cm<sup>3</sup>      D 250 cm<sup>3</sup>

- 18 Which mass of urea, CO(NH<sub>2</sub>)<sub>2</sub>, contains the same mass of nitrogen as 101.1g of potassium nitrate?

- A 22g      B 30g      C 44g      D 60g

- 19 Anhydrous magnesium nitrate, Mg(NO<sub>3</sub>)<sub>2</sub>, will decompose when heated, giving a white solid and a mixture of two gases X and Y.

Y is oxygen.

What is the ratio  $\frac{\text{mass of X released}}{\text{mass of Y released}}$ ?

- A  $\frac{1}{0.174}$       B  $\frac{1}{0.267}$       C  $\frac{1}{0.348}$       D  $\frac{1}{3.43}$



$\frac{92}{16} = \frac{5.75}{1} \times \frac{1/5.75}{1/5.75} = \frac{1}{0.174}$

16.

$$18.5 \text{ mmol/l} \longrightarrow ? \text{ mg/dl}$$

$$\begin{aligned} \text{Mr } \text{C}_6\text{H}_{12}\text{O}_6 &= (12 + 16)6 + 12 \\ &= 180 \end{aligned}$$

$$\begin{aligned} \text{mmol} &= n \times \text{Mr} \\ &= 18.5 \text{ m} \times 180 \\ &= 3330 \text{ mg} \end{aligned}$$

$$3330 \text{ mg/l}$$

$$\begin{array}{ccc} \text{l} & : & \text{dl} \\ 1 & : & 0.1 \\ 3330 & : & x \end{array}$$

$$x = 333 \text{ mol}$$

$$\text{Answer} = 333 \text{ mg/dl}$$

17.



Ratio	1	3	1	1	Liquid
Initial	10	60	-	-	
Reacted	10	30	10	10	
Left over	-	30	10	10	