

Chapter 3 Trigonometry

May/June 2002

1 Prove the identity

$$\cot \theta - \tan \theta \equiv 2 \cot 2\theta. \quad [3]$$

Oct/Nov 2002

5 (i) Express $4 \sin \theta - 3 \cos \theta$ in the form $R \sin(\theta - \alpha)$, where $R > 0$ and $0^\circ < \alpha < 90^\circ$, stating the value of α correct to 2 decimal places. [3]

Hence

(ii) solve the equation

$$4 \sin \theta - 3 \cos \theta = 2,$$

giving all values of θ such that $0^\circ < \theta < 360^\circ$, [4]

(iii) write down the greatest value of $\frac{1}{4 \sin \theta - 3 \cos \theta + 6}$. [1]

May/June 2003

1 (i) Show that the equation

$$\sin(x - 60^\circ) - \cos(30^\circ - x) = 1$$

can be written in the form $\cos x = k$, where k is a constant. [2]

(ii) Hence solve the equation, for $0^\circ < x < 180^\circ$. [2]

10 (i) Prove the identity

$$\cot x - \cot 2x \equiv \operatorname{cosec} 2x. \quad [3]$$

(ii) Show that $\int_{\frac{1}{6}\pi}^{\frac{1}{4}\pi} \cot x \, dx = \frac{1}{2} \ln 2$. [3]

(iii) Find the exact value of $\int_{\frac{1}{6}\pi}^{\frac{1}{4}\pi} \operatorname{cosec} 2x \, dx$, giving your answer in the form $a \ln b$. [4]

Oct/Nov 2003

3 Solve the equation

$$\cos \theta + 3 \cos 2\theta = 2,$$

giving all solutions in the interval $0^\circ \leq \theta \leq 180^\circ$. [5]

May/June 2004

1 Sketch the graph of $y = \sec x$, for $0 \leq x \leq 2\pi$. [3]

5 (i) Prove the identity

$$\sin^2 \theta \cos^2 \theta \equiv \frac{1}{8}(1 - \cos 4\theta).$$

[3]

(ii) Hence find the exact value of

$$\int_0^{\frac{1}{3}\pi} \sin^2 \theta \cos^2 \theta \, d\theta.$$

[3]

Oct/Nov 2004

4 (i) Show that the equation

$$\tan(45^\circ + x) = 2 \tan(45^\circ - x)$$

can be written in the form

$$\tan^2 x - 6 \tan x + 1 = 0.$$

[4]

(ii) Hence solve the equation $\tan(45^\circ + x) = 2 \tan(45^\circ - x)$, for $0^\circ < x < 90^\circ$.

[3]

May/June 2005

6 (i) Prove the identity

$$\cos 4\theta + 4 \cos 2\theta \equiv 8 \cos^4 \theta - 3.$$

[4]

(ii) Hence solve the equation

$$\cos 4\theta + 4 \cos 2\theta = 2,$$

for $0^\circ \leq \theta \leq 360^\circ$.

[4]

Oct/Nov 2005

5 By expressing $8 \sin \theta - 6 \cos \theta$ in the form $R \sin(\theta - \alpha)$, solve the equation

$$8 \sin \theta - 6 \cos \theta = 7,$$

for $0^\circ \leq \theta \leq 360^\circ$.

[7]

May/June 2006

4 (i) Express $7 \cos \theta + 24 \sin \theta$ in the form $R \cos(\theta - \alpha)$, where $R > 0$ and $0^\circ < \alpha < 90^\circ$, giving the exact value of R and the value of α correct to 2 decimal places.

[3]

(ii) Hence solve the equation

$$7 \cos \theta + 24 \sin \theta = 15,$$

giving all solutions in the interval $0^\circ \leq \theta \leq 360^\circ$.

[4]

Oct/Nov 2006

2 Solve the equation

$$\tan x \tan 2x = 1,$$

giving all solutions in the interval $0^\circ < x < 180^\circ$.

[4]

May/June 2007

- 5 (i) Express $\cos \theta + (\sqrt{3}) \sin \theta$ in the form $R \cos(\theta - \alpha)$, where $R > 0$ and $0 < \alpha < \frac{1}{2}\pi$, giving the exact values of R and α . [3]

(ii) Hence show that
$$\int_0^{\frac{1}{2}\pi} \frac{1}{(\cos \theta + (\sqrt{3}) \sin \theta)^2} d\theta = \frac{1}{\sqrt{3}}.$$
 [4]

Oct/Nov 2007

- 5 (i) Show that the equation

$$\tan(45^\circ + x) - \tan x = 2$$

can be written in the form

$$\tan^2 x + 2 \tan x - 1 = 0. \quad [3]$$

- (ii) Hence solve the equation

$$\tan(45^\circ + x) - \tan x = 2,$$

giving all solutions in the interval $0^\circ \leq x \leq 180^\circ$. [4]

May/June 2008

- 4 (i) Show that the equation $\tan(30^\circ + \theta) = 2 \tan(60^\circ - \theta)$ can be written in the form

$$\tan^2 \theta + (6\sqrt{3}) \tan \theta - 5 = 0. \quad [4]$$

- (ii) Hence, or otherwise, solve the equation

$$\tan(30^\circ + \theta) = 2 \tan(60^\circ - \theta),$$

for $0^\circ \leq \theta \leq 180^\circ$. [3]

Oct/Nov 2008

- 6 (i) Express $5 \sin x + 12 \cos x$ in the form $R \sin(x + \alpha)$, where $R > 0$ and $0^\circ < \alpha < 90^\circ$, giving the value of α correct to 2 decimal places. [3]

- (ii) Hence solve the equation

$$5 \sin 2\theta + 12 \cos 2\theta = 11,$$

giving all solutions in the interval $0^\circ < \theta < 180^\circ$. [5]

May/June 2009

- 3 (i) Prove the identity $\operatorname{cosec} 2\theta + \cot 2\theta \equiv \cot \theta$. [3]

- (ii) Hence solve the equation $\operatorname{cosec} 2\theta + \cot 2\theta = 2$, for $0^\circ \leq \theta \leq 360^\circ$. [2]

Oct/Nov 2009/31

5 (i) Prove the identity $\cos 4\theta - 4 \cos 2\theta + 3 \equiv 8 \sin^4 \theta$. [4]

(ii) Using this result find, in simplified form, the exact value of

$$\int_{\frac{1}{6}\pi}^{\frac{1}{3}\pi} \sin^4 \theta \, d\theta. \quad [4]$$

Oct/Nov 2009/32

4 The angles α and β lie in the interval $0^\circ < x < 180^\circ$, and are such that

$$\tan \alpha = 2 \tan \beta \quad \text{and} \quad \tan(\alpha + \beta) = 3.$$

Find the possible values of α and β . [6]

May/June 2010/31

2 Solve the equation

$$\sin \theta = 2 \cos 2\theta + 1,$$

giving all solutions in the interval $0^\circ \leq \theta \leq 360^\circ$. [6]

4 (i) Using the expansions of $\cos(3x - x)$ and $\cos(3x + x)$, prove that

$$\frac{1}{2}(\cos 2x - \cos 4x) \equiv \sin 3x \sin x. \quad [3]$$

(ii) Hence show that

$$\int_{\frac{1}{6}\pi}^{\frac{1}{3}\pi} \sin 3x \sin x \, dx = \frac{1}{8}\sqrt{3}. \quad [3]$$

May/June 2010/32

3 It is given that $\cos a = \frac{3}{5}$, where $0^\circ < a < 90^\circ$. Showing your working and without using a calculator to evaluate a ,

(i) find the exact value of $\sin(a - 30^\circ)$, [3]

(ii) find the exact value of $\tan 2a$, and hence find the exact value of $\tan 3a$. [4]

May/June 2010/33

3 Solve the equation

$$\tan(45^\circ - x) = 2 \tan x,$$

giving all solutions in the interval $0^\circ < x < 180^\circ$. [5]

7 (i) Prove the identity $\cos 3\theta \equiv 4 \cos^3 \theta - 3 \cos \theta$. [4]

(ii) Using this result, find the exact value of

$$\int_{\frac{1}{3}\pi}^{\frac{1}{2}\pi} \cos^3 \theta \, d\theta. \quad [4]$$