

Q1.

3 Fig. 3.1 is a diagram that shows the events that occur between two neurones at a synapse.




Fig. 3.1

(a) Name A to E.

- A
- B
- C
- D
- E

[5]

(b) Draw a large arrow on the diagram to indicate the direction of the impulse across the synapse. [1]

(c) Describe the role of calcium ions in synaptic transmission.


-
-
-
-
-
-

[3]

[Total : 9]

Q2.

- 3 Fig. 3.1 is a diagram of a section through the proximal convoluted tubule of a kidney nephron showing details of cell structure, as seen with the electron microscope.



- (a) Name the structures A and B.

A

B[2]

(b) Explain three ways in which the cells of the proximal convoluted tubule are adapted for selective reabsorption.

1.

.....

2.

.....

3.

.....[3]

(c) Describe the mechanism of glucose reabsorption into the blood from the lumen of the proximal convoluted tubule of the kidney.

.....

.....

.....

.....

.....[3]

(d) Outline, in terms of water potential, how water is reabsorbed by the cells of the proximal convoluted tubule.

.....

.....

.....[2]

[Total: 10]

Q3.

3 Fig. 3.1 is a diagram of a reflex arc.




Fig. 3.1

(a) Explain **briefly** how the stimulus at the finger produces an impulse in the sensory neurone.

.....
.....
.....
.....[3]

(b) Describe the role of the motor neurone in the reflex arc.

.....
.....
.....
.....[3]

(c) Suggest why nerve impulses can only travel in one direction through the reflex arc.

.....
.....
.....[2]

[Total: 8]

Q4.

- 3 During the process of the excretion of nitrogenous waste in mammals, blood passes from the renal artery into networks of capillaries called glomeruli.
Fig. 3.1 is an electronmicrograph showing the relationship between the capillaries and the renal capsule cells, called podocytes.




Fig. 3.1

- (a) Name structures A and B.

A

B

[2]

- (b) Draw an arrow, on Fig. 3.1, to show the passage of fluid out of the capillary.

[2]

- (c) (i) Name the fluid that collects in the capsular space.

.....
.....[1]

- (ii) Describe how the composition of this fluid differs from blood plasma.

.....
.....
.....
.....[2]

- (d) Ultrafiltration involves the removal of small molecules, including urea, from the blood into the renal capsule. Explain what is required for ultrafiltration to occur.

.....
.....
.....
.....
.....
.....
.....

[3]

[Total: 10]

Q5.

- 4 (a) Name the transmitter which is responsible for the transmission of nerve impulses across a cholinergic synapse.

..... [1]

- (b) Outline the role of calcium ions in synaptic transmission.

.....
.....
.....
.....
.....
.....
.....
.....
.....
.....

[4]

- (c) Explain how a synapse ensures one-way transmission of nerve impulses.

.....
.....
.....
.....

[2]

[Total: 7]

Q6.

- 4 (a) Describe the role of insulin in the regulation of blood glucose concentration.

Use

.....
.....
.....
.....
.....
.....
.....
.....[4]

Q7.

- 2 Fig. 2.1 is a light micrograph of a small part of the pancreas.




Fig. 2.1

(a) Name structures **A** and **B**.

A

B [2]

(b) With reference to Fig. 2.1, explain why the pancreas is an *endocrine gland*.

.....

.....

..... [2]

(c) Insulin and glucagon are involved in the control of blood glucose concentration. When blood glucose concentration rises, secretion of insulin increases.

Outline two ways in which insulin affects the activity of cells in the liver.

1.

.....

2.

..... [2]

(d) Some forms of diabetes are caused by an inability to secrete insulin and can be controlled by regular injections of insulin. Most of this insulin is now produced using genetically modified *Escherichia coli*.

Explain the advantages of using this type of insulin, rather than insulin obtained from animal sources.

.....

.....

.....

..... [2]

[Total: 8]

A
Exam
U

Q8.

4 Almost 40% of adults with cystic fibrosis (CF) develop a form of diabetes known as cystic fibrosis-related diabetes (CFRD). This is thought to happen because the build-up of thick secretions in the pancreas destroys β cells.

For
Examin
Use

(a) Explain how the destruction of β cells causes diabetes.

.....

.....

.....

.....

.....

.....

.....

.....

.....

[4]

(b) The bacterium *Pseudomonas aeruginosa* can cause chronic (long-lasting) lung infections. A person with CFRD is likely to have poorer lung function and a greater likelihood of having a chronic lung infection than a person who has CF but does **not** have CFRD.

An investigation was carried out to find out if the severity of damage to lung function in a person with CFRD is affected by

- their gender
- whether or not they have a chronic *P. aeruginosa* infection.

The investigators measured lung function by recording the maximum volume of air that can be expelled from the lungs in the first one second of a forced expiration. This is known as FEV₁. The lower the median FEV₁, the poorer the lung function.

	without chronic <i>P. aeruginosa</i> infection				with chronic <i>P. aeruginosa</i> infection			
	male		female		male		female	
	with CFRD	without CFRD	with CFRD	without CFRD	with CFRD	without CFRD	with CFRD	without CFRD
number of people	44	110	52	93	106	166	121	120
FEV ₁	71.1	71.4	53.6	73.6	49.0	59.0	42.0	61.0

With reference to Table 4.1

- (i) discuss whether or not there appears to be a positive correlation between having a chronic *P. aeruginosa* infection and having CFRD

.....

.....

.....

.....

.....

.....

.....

[2]

- (ii) calculate the percentage difference between the FEV₁ of males and females without CFRD and without *P. aeruginosa* infection.

Show your working

answer % [2]

- (iii) outline the conclusions that can be drawn concerning the relationship between gender and the severity of lung damage in a person with CFRD and with *P. aeruginosa* infection. Ex

.....

.....

.....

.....

.....

.....

.....

.....

.....

..... [3]

- (c) In a person with CF, damage to lung function and the increased likelihood of chronic infections are the result of the secretion of thick mucus.

Explain why thick mucus is secreted in the lungs of a person with CF.

.....

.....

.....

.....

.....

.....

.....

.....

.....

..... [4]

[Total: 15]

Q9.

6 (a) Fig. 6.1 outlines how a cholinergic synapse works.




Fig. 6.1

With reference to Fig. 6.1:

(i) name **A** and **B**

A

B [2]

(ii) name the process occurring at **C**

..... [1]

(iii) state the effect of **B** entering the post-synaptic neurone

..... [1]

(iv) explain the role of acetylcholinesterase in the synapse.

.....

.....

.....

.....

.....

..... [3]

(b) Some synapses in the brain use the neurotransmitter dopamine. After the postsynaptic membrane has been depolarised, dopamine leaves the receptor proteins and moves back into the presynaptic neurone through specific transporter proteins.

Schizophrenia is a condition in which there is a higher than usual concentration of dopamine in certain areas of the brain. The drug phenothiazine has a similar shape to dopamine and is used to treat schizophrenia.

Suggest and explain what occurs at the synapse when phenothiazine is used in the treatment of schizophrenia.

.....

.....

.....

..... [2]

Ex

Q10.

6 (a) The human kidneys process 1200 cm^3 of blood every minute. This 1200 cm^3 of blood contains 700 cm^3 of plasma. As blood passes through the glomeruli of the kidneys, 125 cm^3 of fluid passes into the renal capsules (Bowman's capsules). This fluid is called the glomerular filtrate and is produced by a process called ultrafiltration.

For
Examin
Use

(i) Calculate the percentage of plasma that passes into the renal capsules.

Show your working and **give your answer to one decimal place.**

answer% [2]

(ii) Explain how the **structures** of the glomerular capillaries and the podocytes are adapted for ultrafiltration.

glomerular capillaries

.....
.....
.....
.....

podocytes

.....
.....
.....
.....

[4]

- (b) The glomerular filtrate then passes through the proximal convoluted tubule.

Fig. 6.1 is a transverse section through part of the proximal convoluted tubule.




Fig. 6.1

- (i) Name the structures labelled X.

..... [1]

- (ii) Explain why the epithelial cells of the proximal convoluted tubule have many mitochondria in them.

.....
.....
.....
..... [2]

- (iii) Of the 125cm^3 of glomerular filtrate that enters the renal capsules each minute, only 45cm^3 reaches the loops of Henlé.

Name **two** substances that are reabsorbed into the blood from the proximal convoluted tubule, **apart from water**.


.....
..... [2]

[Total: 11]

Q11.

- 3 The mammalian liver is made up of lobules that consist of liver cells (hepatocytes) arranged in plates. Between these plates of cells are enlarged leaky capillaries called sinusoids. Blood from both the hepatic portal vein and the hepatic artery flows through these sinusoids to the central vein and eventually into the hepatic vein. Inside the sinusoids are Kupffer cells. Fig. 3.1 shows a section of a liver lobule and its associated blood vessels.

Use



- (a) Describe the role of the Kupffer cells in the homeostatic function of the liver.

.....
.....
.....
.....
.....[4]

(b) State how liver cells are involved in fat metabolism.

.....
.....
.....
.....[3]

(c) Explain how urea produced by liver cells from the deamination of excess amino acids is transported to the kidney for excretion.

.....
.....
.....[2]

(d) State how blood in the hepatic vein will differ after a heavy meal from blood in

(i) the hepatic portal vein;

.....
.....[1]

(ii) the hepatic artery.

.....
.....[1]

[Total : 11]

Q12.

3 Figs 3.1 and 3.2 show the concentration of glucose and insulin in blood plasma before and after a glucose drink.




Fig. 3.1




Fig. 3.2

(a) With reference to Fig. 3.1, describe the changes in blood glucose concentration after the glucose drink.

.....

.....

.....

.....

[3]

(b) With reference to Fig. 3.1 and Fig. 3.2, explain how the changes in blood glucose cause:

(i) an increase in the concentration of insulin in the plasma;

.....

.....

.....

[2]

(ii) a subsequent fall in the concentration of insulin in the plasma.

.....

.....

.....

[2]

(c) Describe the role of the hormone glucagon in maintaining the concentration of blood glucose.

.....
.....
.....
.....[3]

[Total : 10]

Q13.

5 Fig. 5.1 is a drawing of a section of a liver lobule that has been injected with ink. The Kupffer cells are clearly visible as a result of taking up carbon particles from the ink by phagocytosis.




Fig. 5.1

(a) The Kupffer cells remove damaged red blood cells from the blood in the sinusoids. Explain what happens to the haemoglobin.

.....
.....
.....
.....
.....
.....[4]

(b) Describe how excess amino acids are deaminated by the hepatocytes.

.....
.....
.....
.....
.....[3]

(c) Outline the function of the hepatocytes in detoxification of a **named** toxic compound.

.....
.....
.....
.....
.....[3]

[Total : 10]

Q14.

- 2 Fig. 2.1 shows the changes in membrane potential in an axon during the passage of a single impulse.




Fig. 2.1

- (a) Outline how the resting potential from A to B is maintained.

.....
.....
.....
.....
.....
.....[3]

- (b) Describe how the changes in the membrane bring about depolarization from B to C.

.....
.....
.....
.....
.....
.....[3]

(c) Explain how the membrane is repolarised from C to D.

.....
.....
.....
.....
.....[3]

(d) State three differences between nervous and hormonal communication in mammals.

1.
.....
2.
.....
3.
.....
[3]

[Total : 12]

Q15.

3 In mammalian kidneys, the loop of Henle is closely associated with the process of osmoregulation.

(a) Explain what is meant by osmoregulation.

.....
.....
.....
.....[2]

Fig. 3.1 shows the water potential of renal fluid as it passes through the loop of Henle.




Fig. 3.1

- (b) Using the information given in Fig. 3.1, describe and explain what happens to the renal fluid as it passes through the loop of Henle.

.....
.....
.....
.....
.....
.....
.....
.....
..... [5]

- (c) Control systems often work by using negative feedback. These systems require a receptor and an effector. In the process of osmoregulation name the receptor and effector involved.

Receptor

Effector

[2]

[Total: 9]

Q16.

- 5 The lower epidermis of a dicotyledonous leaf is perforated with stomata. Each stoma is bounded by two guard cells which control the size of the pore.

Fig. 5.1 shows a scanning electron microscope photograph of an open and a closed stoma.




Fig. 5.1

X4000

- (a) Calculate the length of the pore of the open stoma in Fig. 5.1.

Show your working.

Answer

[2]

- (b) Explain the importance of abscisic acid, ABA, in causing stomatal closure.

.....
.....
.....
.....
.....
.....
.....
.....
.....
.....

[4]

- (c) Under conditions of low wind speed, the rate of transpiration decreases, even though the stomata of the leaves are open.

Explain why this is so.

.....
.....
.....
.....

[2]

[Total: 8]

Q17.

4 (a) Explain the role of negative feedback in homeostasis in mammals.

.....
.....
.....
.....
.....
.....
.....
..... [4]

(b) The enzyme glucose oxidase catalyses the conversion of glucose to gluconic acid.



Describe how glucose oxidase in a biosensor can give warning of low blood glucose concentration (hypoglycaemia).

.....
.....
.....
.....
.....
.....
.....
..... [4]

[Total: 8]

Q18.

- 8 Fig. 8.1 shows the changes in potential difference (p.d.) across the membrane of a neurone over a period of time. The membrane was stimulated at time A and time B with stimuli of different intensities.




Fig. 8.1

- (a) Stimulus B resulted in an action potential. Describe what is occurring at C, D and E.

C

.....

.....

.....

D

.....

.....

.....

E

.....

.....

.....

[6]

(b) Suggest why stimulus **A** did not result in an action potential being produced whereas stimulus **B** did.

.....
.....
.....
.....
.....
.....

[2]

[Total: 8]

Q19.

7 Fig. 7.1 shows a section through part of the cortex of a kidney.




Fig. 7.1

(a) On Fig. 7.1, draw label lines and use the letters **G** and **R** to identify :

- a glomerulus with the letter **G**.
- a renal capsule with the letter **R**.

[2]

(b) State the name of the hormone that is involved in the control of the water potential of the blood.

.....[1]

(c) Table 7.1 shows the concentration of some compounds in the fluids of a glomerulus, a renal capsule and a collecting duct of the kidney.

Ex

Table 7.1

compound	concentration / g 100 cm ⁻³		
	blood plasma entering glomerulus	filtrate in renal capsule	urine in collecting duct
water	90	90	96
proteins	8.0	0.0	0.0
glucose	0.1	0.1	0.0
urea	0.03	0.03	2.0

With reference to Table 7.1,

(i) explain why proteins occur in the blood entering the glomerulus but not in the filtrate in the renal capsule

.....
.....
.....[2]

(ii) explain why there is glucose present in the filtrate but not in the urine

.....
.....
..... [2]

(iii) explain the difference in the concentration of urea between the filtrate and urine.

.....
.....
..... [2]

[Total: 9]

Q20.

6 (a) The pancreas acts both as an exocrine and an endocrine gland.

(i) Describe the parts of the pancreas involved in its **endocrine** function.

.....
.....
.....
.....
.....
..... [3]

(ii) State precisely the group of compounds to which the pancreatic hormone insulin belongs.

.....[1]

(b) People with insulin-dependent (type 1) diabetes require regular injections of insulin. In the past the insulin used came from animal sources such as pigs. Diabetics now use human insulin that has been manufactured using gene technology.

Describe the advantages of treating diabetics with insulin produced by gene technology.

.....
.....
.....
.....
.....
.....
.....[3]

[Total: 7]

Q21.


7 An investigation was carried out into the effects of a plant growth regulator, auxin (IAA), on apical dominance.

- The apical buds of 20 pea plants were cut off and discarded.
- The cut surfaces of 10 pea plants were coated with an inert paste containing auxin.
- The cut surfaces of the other group of 10 pea plants were coated with the inert paste alone.
- A further group of 10 pea plants did not have their apical buds removed and were not coated with paste. This was a control group.

The lengths of the side shoots of plants in each of the three groups were measured at regular time intervals and mean values calculated.

The results are shown in Fig. 7.1.

Exa
t



(a) Explain why the side shoots increase in length when the terminal buds are removed.

.....
.....
.....
.....
.....
..... [3]

f
Exam
L

(b) Calculate the percentage difference, at 13 days, in the mean length of side shoots of plants treated with paste alone compared with the plants treated with paste and auxin.

Give your answer to the nearest whole number.

Show your working.

Answer % [2]

(c) Using data from Fig. 7.1, describe **and** explain the effect of auxin on the growth of side shoots.

.....
.....
.....
.....
.....
..... [3]

[Total: 8]

Q22.

7 Urea is the main nitrogenous waste product in humans. It is made in the liver and excreted by the kidneys in urine.

B

(a) Define the term *excretion*.

.....
.....
.....
..... [2]

(b) The kidneys regulate the water potential of body fluids. This is known as osmoregulation and involves a negative feedback system.

Outline the role of negative feedback in osmoregulation.

.....
.....
.....
.....
.....
.....
.....
..... [4]

(c) Investigations have shown that when a person remains in a cold environment for more than 15 minutes there is increased urine production. This is called cold diuresis.

Suggest how exposure to cold can lead to cold diuresis.

.....
.....
.....
..... [2]

[Total: 8]

Q23.

- 2 (a) The first diagnostic test strip using immobilised enzymes was a dip stick to estimate the concentration of glucose in urine.

The dip stick is a thin strip of plastic with a cellulose pad containing two enzymes and a colour reagent (chromogen) at one end. The pad responds with a colour change after being dipped into a sample of urine that contains glucose. The colour can be matched against a graded colour chart to give a 'semi-quantitative' estimate of the concentration of glucose in the sample, as shown in Fig. 2.1. The chart shows the colours of a negative reaction (–) and three increasingly positive reactions (+, ++ and +++).




Fig. 2.1

Explain why the estimate of glucose concentration achieved by this method is only 'semi-quantitative'.

.....


.....

.....

.....

..... [2]

- (b) One of the two enzymes immobilised in the cellulose pad on the test strip is glucose oxidase, which catalyses the following reaction:



This reaction does **not** result in the development of colour by the chromogen. This is achieved by the activity of the second immobilised enzyme in the pad.

- (i) Name the second immobilised enzyme in the pad.

.....[1]

- (ii) Explain how the reaction catalysed by this enzyme results in the chromogen changing colour.

.....
.....
.....
.....[2]

- (iii) The cellulose pad on the test strip is covered by a layer of cellulose acetate, which is permeable to glucose molecules, but not to larger molecules.


Suggest why the layer of cellulose acetate is present.

.....
.....
.....
.....[2]

Exa
t

- (c) Two young men, subjects **A** and **B**, were each given a standardised test dose of glucose after fasting.
- The blood glucose concentration of each subject was then measured immediately and at 30 minute intervals for two hours.
 - Samples of their urine were taken and tested at the same time intervals. The colour change of each test strip was compared with the colour chart and recorded as **-**, **+**, **++** or **+++**.

The results of the investigation are shown in Fig. 2.2.



Results of urine tests:

	time after test dose of glucose given/h				
subject	0.0	0.5	1.0	1.5	2.0
A	-	-	-	-	-
B	-	+	++	++	+++

Fig. 2.2

With reference to Fig. 2.2:

Ex

- (i) explain the differences between the **blood glucose** concentrations of **A** and **B**

.....
.....
.....
.....
.....
.....
.....
.....
.....
..... [4]

- (ii) suggest what is meant by the term 'renal threshold'

.....
..... [1]

- (iii) describe the events in the kidneys, after ultrafiltration, that result in the increasing quantity of glucose in **B**'s urine.

.....
.....
.....
.....
.....
.....
..... [3]

[Total: 15]

Q24.

Fig. 9.2 shows synapses in a pain pathway from a damaged joint to the brain.




Fig. 9.2

E

Table 1.1

event	description of event
A	Calcium ions enter presynaptic neurone knob.
B	Acetylcholine binds to receptor proteins on postsynaptic membrane.
C	Vesicles fuse with presynaptic membrane and release acetylcholine into synaptic cleft.
D	Postsynaptic membrane becomes depolarised.
E	Nerve impulse reaches presynaptic membrane.
F	Acetylcholine diffuses across cleft.
G	Receptor proteins change shape, channels open and sodium ions enter postsynaptic neurone.
H	Calcium ion channels open in presynaptic membrane.
I	Nerve impulse generated in postsynaptic neurone.
J	Vesicles of acetylcholine move towards presynaptic membrane.

Complete Table 1.2 to show the events in the correct order.

Two of the events have been done for you.

Table 1.2

correct order	letter of stage
1	E
2	
3	
4	
5	
6	F
7	
8	
9	
10	

[4]

(b) Synapses have many roles in nervous coordination in mammals.

(i) Explain how synapses ensure one-way transmission of nerve impulses.

.....
.....
.....
.....
.....
.....
.....

[2]

- (ii) In a learning activity, it is believed that the number of synapses between brain neurones increases.
Suggest the advantages of this increased number of synapses.

Ex

.....
.....
.....
.....
.....
.....

[2]

[Total: 8]

Q27.

- 1 (a) Fig. 1.1 shows a neurone forming three synapses with adjacent neurones.




Fig. 1.1

Name A, B and C.

A.....

B.....

C.....

[3]

(b) Outline the role of structure **A** in synaptic transmission.

.....
.....
.....
.....
.....
.....
.....
.....
.....
..... [3]

(c) The drug nicotine has a similar structure to acetylcholine.

Suggest the effects on brain neurones of inhaling nicotine from a cigarette.

.....
.....
.....
.....
.....
.....
..... [2]

[Total: 8]

Q28.

- 6 (a) Table 6.1 shows the mean axon diameter and mean speed of conduction of nerve impulses for four different animals.

Table 6.1

animal	type of neurone	axon diameter / μm	mean speed of conduction / ms^{-1}
A – mammal	myelinated	4	25
B – mammal	unmyelinated	5	3
C – amphibian	myelinated	14	35
D – amphibian	myelinated	10	30

With reference to Table 6.1, describe:

- (i) the effect of myelination on the speed of conduction of impulses in mammals

.....
.....
.....
..... [2]

- (ii) the effect of axon diameter on the speed of conduction of impulses in amphibians.

.....
.....
.....
..... [2]

- (b) Explain how myelination affects the speed of conduction of impulses.

.....
.....
.....
.....
.....
.....
.....
..... [3]

(c) Multiple sclerosis (MS) is an auto-immune condition of humans in which the body's immune system attacks the myelin sheaths which are then damaged. This leads to a decrease in information reaching the brain from sensory receptors. E

(i) Suggest how the myelin sheaths may be attacked.

.....
.....
.....
.....[2]

(ii) Explain why this damage leads to a decrease in information reaching the brain from sensory receptors.

.....
.....
.....
.....[2]

[Total: 11]

Q29.

- 9 The passage below summarises the effects of gibberellins on seed germination.

Complete the passage by using the most appropriate scientific term(s).

When a seed is shed from the parent plant, it is in a state of ,
which means it is metabolically inactive.

When water is absorbed by a seed, it stimulates the production of gibberellin by the
..... within the seed. The gibberellin stimulates the synthesis of
amylase by cells in the layer.

Amylase hydrolyses starch molecules in the converting them
to soluble molecules. These molecules are converted to
glucose which is transported to the embryo, providing a source of carbohydrate that can be
respired to provide as the embryo begins to grow.

Gibberellin causes these effects by regulating genes that are involved in the synthesis of
amylase. It has been shown that application of gibberellin to seeds can cause an increase in
the of the DNA coding for amylase.

[Total: 7]

Q30.

- 6 Fig. 6.1 is a trace that shows the changes that occur in the membrane potential of a neurone during an action potential.




Fig. 6.1

- (a) Using the letter(s) **A** to **F** from Fig. 6.1, state which letter(s) corresponds to the following:

- (i) depolarisation
- (ii) hyperpolarisation
- (iii) the membrane is most permeable to potassium ions
- (iv) resting potential [4]

- (b) Saxitoxin is a powerful poison produced naturally by single-celled, eukaryotic, photosynthetic, marine organisms. Shellfish may consume organisms containing saxitoxin but are unaffected. If humans were to eat shellfish containing saxitoxin they would become very ill and may die.

- (i) State the kingdom to which the organisms that produce saxitoxin belong.

.....[1]

- (ii) Saxitoxin blocks sodium ion channels in the cell surface membranes of neurones.
Describe the role of sodium ion channels in the transmission of a nerve impulse.

B

.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
..... [3]

- (iii) Suggest why saxitoxin may be fatal to humans.

.....
.....
.....
..... [2]

[Total: 10]

Q31.

- 9 The passage below summarises the effects of auxin on the growth of a shoot.
Complete the passage by using the most appropriate scientific term(s).

1
Exam
1

Auxin is synthesised in the growing tips of shoots (apical buds). It is transported from here down the shoot by from cell to cell and also to a lesser extent by flow in the

Auxin seems to be involved in determining whether a plant grows upwards or whether it branches sideways. When the apical bud is actively growing, it tends to stop lateral buds from growing. This is called apical The plant grows upwards rather than branching out sideways.

However, if the apical bud is cut off, the lateral buds start to grow. It is thought that removal of the apical bud causes the concentration of auxin in lateral buds to so the buds can now grow by cell and cell

[Total: 7]

Q32.

- 6 (a) The passage below outlines how sensory receptors work.

Complete the passage by using the most appropriate scientific term(s).

A sensory receptor cell responds to a stimulus by opening ion in its cell surface membrane. Sodium ions flood into the cell causing the membrane to become This is called the potential. If this potential is large enough to reach a then an action potential is transmitted to the central nervous system. An increase in the strength of the stimulus will result in an increase in the of action potentials transmitted. [5]

6 (a) In mammals, the water potential of the blood is constantly monitored by osmoreceptor cells in the hypothalamus of the brain. When the water potential of the blood decreases, ADH (antidiuretic hormone) is produced by cells in the hypothalamus and released into the blood via an endocrine gland.

(i) Explain what is meant by the term *water potential*.

.....
.....
.....[1]

(ii) Describe the effect on water potential of adding solute to a solution.

.....
.....[1]

(iii) State precisely where ADH is released into the blood.

.....[1]

(iv) The decrease in the water potential of the blood is sometimes due to the loss of water from the body of a mammal.

List two ways by which water may be lost from the body.

.....
.....[1]

6 (a) Neurones transmit impulses from one part of a mammal's body to another.

The table contains statements that refer to motor and sensory neurones.

Complete the table, indicating with the letters **M**, **S** or **B**, whether each statement applies to:

- motor neurones only (**M**)
- sensory neurones only (**S**)
- both motor and sensory neurones (**B**).

The first one has been done for you.

statement	letter
is myelinated	B
may form a synapse with an intermediate (relay) neurone
cell body lies within the CNS
dendron is usually longer than axon
cell body lies within spinal nerve
has many dendrites

[3]

6 Asparagus, *Asparagus officinalis*, is a green vegetable plant grown in many parts of the world.

Fig. 6.1 shows some asparagus shoots.



Fig. 6.1

(a) The shoots are usually cooked and then eaten hot or cold. Asparagus contains many mineral ions which have important roles in the bodies of mammals.

Complete Table 6.1 to summarise some of these roles.

Table 6.1

ion	role	type of cell
Fe^{2+}	red blood cell
Na^+	co-transport in the kidney
Ca^+	neurone

[3]

- 6 (a) Explain how a synapse functions. [9]
(b) Describe the role of glucagon in regulating blood glucose. [6]
[Total: 15]

2.

- 6 (a) Describe the role of auxins in apical dominance. [6]
(b) Explain the role of gibberellins in the germination of wheat or barley. [9]
[Total: 15]

3.

- 6 (a) Explain the meaning of the term **homeostasis** with specific reference to the control of **raised** blood glucose concentration in mammals. [8]
(b) Describe the role played by ADH in osmoregulation in mammals. [7]
[Total: 15]

4.

- 9 (a) Describe how a nerve impulse crosses a cholinergic synapse. [9]
(b) Explain the roles of synapses in the nervous system. [6]
[Total: 15]

5.

- 10 (a) Compare the roles of the endocrine **and** nervous systems in control and coordination in animals. [8]
(b) Describe the part played by auxins in apical dominance in a plant shoot. [7]
[Total: 15]

6.

- 10 (a) Describe the part played by the proximal convoluted tubules in the functioning of the kidneys. [8]
- (b) Explain how the collecting ducts in the kidneys may reduce the loss of water from the body. [7]
- [Total: 15]

7.

- 10 (a) Describe the structure of a kidney nephron and its associated blood vessels. [7]
- (b) Explain how glomerular filtrate is formed. [8]
- [Total: 15]

8.

- 10 (a) Describe the structure of a myelinated sensory neurone. [7]
- (b) Explain how an action potential is transmitted along a sensory neurone. [8]
- [Total: 15]

9.

- 9 (a) Outline the ways in which the endocrine **and** nervous systems carry out their roles in control and coordination in animals. [8]
- (b) Describe the part played by auxins in apical dominance in a plant shoot. [7]
- [Total: 15]

10.

- 11 (a) Describe the role of abscisic acid (ABA) in the closure of a stoma. [8]
- (b) Describe the role of gibberellins in the germination of barley seeds. [7]
- [Total: 15]

11.

- 10 (a) Describe the structure of a kidney, including its associated blood vessels. [6]
- (b) Describe the mechanisms involved in reabsorption in the proximal convoluted tubule **and** describe how the epithelial cells of the proximal convoluted tubule are adapted to carry out this process. [9]

[Total: 15]

12.

- 10 (a) Outline, with reference to blood glucose concentration, the principles of homeostasis in mammals. [6]
- (b) Describe the roles of the endocrine and nervous systems in control and coordination in mammals. [9]

[Total: 15]

13.

- 7 (a) Describe how nitrogenous waste products are formed and explain why they need to be removed from the body. [6]
- (b) Describe how the kidney removes metabolic wastes from the body. [9]

14.

- 6 (a) Describe how the structure of neurones speeds up the transmission of action potentials. [6]
- (b) Explain, using a named example, how sensory receptors in mammals convert energy into action potentials. [9]

[Total: 15]

15.

- 10 (a) Describe the structure of a motor neurone. [7]
- (b) Explain how an action potential is transmitted along a motor neurone. [8]

[Total: 15]

16.

- 10 (a) Describe a reflex arc **and** explain why such reflex arcs are important. [7]
- (b) Describe the structure of a myelin sheath **and** explain its role in the speed of transmission of a nerve impulse. [8]
- [Total: 15]

17.

- 10 (a) Describe how a nerve impulse crosses a cholinergic synapse. [9]
- (b) Explain the roles of synapses in the nervous system. [6]
- [Total: 15]

18.

- 11 (a) Describe how a resting potential is maintained in an axon. [9]
- (b) Describe, using named examples, how sensory receptors in mammals generate action potentials. [6]
- [Total: 15]

19.

- 10 (a) Outline the process of the photolysis of water **and** describe what happens to the products of photolysis. [10]
- (b) Describe the roles of gibberellins in stem elongation. [5]
- [Total: 15]

20.

- 9 (a) Describe the action of glucagon on liver cells in the regulation of blood glucose concentration. [9]
- (b) Outline how a dip stick can measure the concentration of glucose **and** suggest advantages of using an electronic biosensor instead of a dip stick. [6]
- [Total: 15]

[Online Classes : Megalecture@gmail.com](mailto:Megalecture@gmail.com)
www.youtube.com/megalecture
www.megalecture.com

[Online Classes : Megalecture@gmail.com](mailto:Megalecture@gmail.com)
www.youtube.com/megalecture
www.megalecture.com

[Online Classes : Megalecture@gmail.com](mailto:Megalecture@gmail.com)
www.youtube.com/megalecture
www.megalecture.com

[Online Classes : Megalecture@gmail.com](mailto:Megalecture@gmail.com)
www.youtube.com/megalecture
www.megalecture.com

[Online Classes : Megalecture@gmail.com](mailto:Megalecture@gmail.com)
www.youtube.com/megalecture
www.megalecture.com

[Online Classes : Megalecture@gmail.com](mailto:Megalecture@gmail.com)
www.youtube.com/megalecture
www.megalecture.com