

5 Chemical energetics

Enthalpy changes and entropy changes accompany chemical reactions. This topic demonstrates why some reactions and processes are spontaneous and others are not.

5.1 Lattice energy



5.2 ionic solid, its aqueous solution and Born-Haber cycles



ENERGETICS

5 Chemical energetics

Enthalpy changes and entropy changes accompany chemical reactions. This topic demonstrates why some reactions and processes are spontaneous and others are not.


Learning outcomes

Candidates should be able to:

- 5.1 Enthalpy change, ΔH**
- explain that chemical reactions are accompanied by energy changes, principally in the form of heat energy; the energy changes can be exothermic (ΔH is negative) or endothermic (ΔH is positive)
 - explain and use the terms:
 - enthalpy change of reaction and standard conditions*, with particular reference to: formation, combustion, hydration, solution, neutralisation, atomisation
 - bond energy* (ΔH positive, i.e. bond breaking)
 - lattice energy*** (ΔH negative, i.e. gaseous ions to solid lattice)
 - calculate enthalpy changes from appropriate experimental results, including use of the relation $q = mc\Delta T$
 - explain the effect of ionic charge and of ionic radii on lattice energy
- 5.2 Hess' Law, including Born-Haber cycles**
- construct and interpret enthalpy changes found by direct experiment, e.g. enthalpy change of formation from enthalpy changes of combustion
 - average bond enthalpies
 - the formation of the ionic solid and of its aqueous solution**
 - Born-Haber cycles including ionisation energy and electron affinity**
 - construct and interpret reaction pathway diagram, in terms of the enthalpy change of reaction and of the activation energy

LATTICE ENERGY

Lattice enthalpy is the energy that would be given out to the surroundings (red arrows) if one mole of a compound could be formed directly from free gaseous ions coming together (black arrows) and arranging themselves into a crystal lattice.



LATTICE ENERGY EXAMPLE

As gaseous ions form ionic solids, energy is released. This energy is known as the lattice enthalpy.



ionic solid → Gaseous ion



Lattice enthalpy is a measure of the strength of the forces between the ions in an ionic solid. The greater the lattice enthalpy, the stronger the forces

LATTICE ENERGY

Lattice energy: the energy change (liberated) when:

1 mole of solid lattice is formed

from its constituent **gaseous** ions

under standard conditions (at 298K and 1 atm).



this value is always negative

because lattice energy is always exothermic

LATTICE ENERGY

Lattice enthalpies are important because they are used as a measure of the strength of ionic bonds in different ionic compounds.

The strengths of ionic bonds, measured as lattice enthalpies (kJ mol^{-1}), arise from the energy given out as billions upon billions of positive and negative ions come together to form a crystal lattice.

The force of attraction between the ions is stronger, and this results in a more exothermic lattice enthalpy, if:


- the charges on the ions are large
- the ionic radii are small, allowing the ions to get closer to each other.

LATTICE ENERGY

The magnitude of Lattice energy depends on the the electrostatic attraction between the ions. It

increases as the charges on the ions increases

decreases as the size of the ions increases.



THE DISTANCE BETWEEN THE IONS

The smaller the distance between the ions, the greater the attraction between them, and the greater will be the lattice enthalpy.

For example, the size of the halide ions increases in the order $\text{F}^- < \text{Cl}^- < \text{Br}^- < \text{I}^-$.

Lattice enthalpies for the sodium halides decreases in the same order (Note that each substance has the same ionic charges).

As the size of an anion increases, attraction between ions decreases and so does the lattice enthalpy.

$$\text{NaF } \Delta H_{\text{lat}} = -926 \text{ kJ mol}^{-1}$$

$$\text{NaCl } \Delta H_{\text{lat}} = -787 \text{ kJ mol}^{-1}$$

$$\text{NaBr } \Delta H_{\text{lat}} = -752 \text{ kJ mol}^{-1}$$

LATTICE ENERGY AND CHARGE ON IONS

The greater the charges on the ions, the greater the attraction between them, and the greater will be the lattice enthalpy.

E.g., sodium fluoride(Na^+F^-) and magnesium oxide($\text{Mg}^{2+}\text{O}^{2-}$) have similar structures.

The sizes of the cations are similar as both Na and Mg are in period 3, and the sizes of the anions are also similar as O and F are in Period 2.

The lattice enthalpy of the oxide is expected to be four times that of the fluoride.

$$\text{NaF } \Delta H (\text{lattice}) = - 926 \text{ kJ mol}^{-1}$$

$$\text{MgO } \Delta H (\text{lattice}) = - 3800 \text{ kJ mol}^{-1}$$

QUESTIONS

1. Explain the term lattice energy. [2]
2. Write a balanced equation including state symbols to represent the lattice energy of magnesium chloride. [2]

QUESTIONS

3. Describe how, and explain why, the lattice enthalpy of magnesium fluoride differs from that of calcium chloride.

4. How would you expect the magnitude of lattice energy of magnesium oxide to compare with that of sodium oxide? Explain your reasoning.

BORN-HABER CYCLE

Born–Haber cycles are an application of Hess's Law. They provide a model for the determination of lattice enthalpies which cannot be measured experimentally. They also enable chemists to test the thermodynamic stability of ionic compounds.


A Born–Haber cycle identifies all the enthalpy changes that contribute to the standard enthalpy change of formation of a compound. These changes involve:

- the energy (enthalpy changes) required to create free gaseous ions by atomising and then ionising the elements
- the energy given out (the lattice enthalpy) when the ions come together to form a crystal lattice.

BORN-HABER CYCLE

A Born-Haber cycle is usually set out like an enthalpy profile diagram, with enthalpy changes one after another.

All the processes in the cycle can be measured experimentally except the lattice enthalpy. So, by using Hess's law it is possible to calculate the lattice enthalpy.



BORN-HABER CYCLE

A method for analysis of the thermochemical cycle based on Hess' law.

With the calculation of lattice enthalpies the following are determined:

atomisation energies for the elements,


ionisation energies for the metal component, ΔH_{ie} ,

electron affinities for the non-metal component, ΔH_{ea} , and

the enthalpy change of formation, ΔH_f , for the compound.

$$\Delta H_{\text{formation}} = \text{Atomisation energy of metal} + \text{Ionisation Energy of Metal} \\ + \text{Atomisation of non-metal} + \text{electron affinity of non-metal} + \text{Lattice energy}$$

GENERAL BORN-HABER CYCLE



FIRST ELECTRON AFFINITY

Electron affinity of an element is the enthalpy change that occurs when one mole of its gaseous atoms gains one mole of electrons under standard conditions.



The first electron affinity is the enthalpy change when 1 mole of electrons is added to 1 mole of gaseous atoms to form 1 mole of gaseous 1^- ions under standard conditions.

Most first electron affinities are exothermic. This is because energy is released when the electron comes under the attraction of the nucleus

Element	Si	P	S	Cl
E.A/ kJ mol^{-1}	-135	-60	-200	-364


SUBSEQUENT ELECTRON AFFINITIES

The subsequent electron affinities are always endothermic since the initial repulsion between the negatively charged ion and the incoming electron outweighs the energy released when the electron comes under the attraction of the nucleus.



The second electron affinity is the enthalpy change when 1 mole of electrons is added to 1 mole of gaseous 1^- ions to form 1 mole of gaseous 2^- ions under standard conditions.

The overall enthalpy change in forming an oxide ion, O^{2-} , from an oxygen atom is found by adding together the 1st and 2nd electron affinities.



EXAMPLE 5

Calculate the lattice energy of sodium chloride using the following data:

Enthalpy of atomisation of sodium = +91 kJ mol⁻¹


Bond Enthalpy of chlorine = +242 kJ mol⁻¹

First ionisation energy of sodium = +496 kJ mol⁻¹

Electron affinity of chlorine = -364 kJ mol⁻¹

Enthalpy of formation of sodium chloride = -411 kJ mol⁻¹

GENERAL BORN-HABER CYCLE



QUESTION 1

Calculate the enthalpy change of formation of potassium fluoride

Enthalpy of atomisation of K = 90 kJ mol⁻¹


Bond energy of fluorine = 158 kJ mol⁻¹

First ionisation energy of K = 420 kJ mol⁻¹

Electron affinity of fluorine = - 333 kJ mol⁻¹

Lattice energy of KF = - 813 kJ mol⁻¹


QUESTION 1



QUESTION 2

Calculate the enthalpy change of formation of calcium chloride

Enthalpy of atomisation of Ca	178 kJ mol ⁻¹
Enthalpy of atomisation of chlorine	121 kJ mol ⁻¹
First ionisation energy of Ca	590 kJ mol ⁻¹
Second energy of ionisation of Ca	1100 kJ mol ⁻¹
Electron affinity of chlorine	- 364 kJ mol ⁻¹
Lattice energy of CaCl ₂	- 2197 kJ mol ⁻¹



QUESTION 3

Calculate the lattice energy of sodium oxide from the following data:

Enthalpy of atomisation of Na	$+91 \text{ kJ mol}^{-1}$
Enthalpy of atomisation of Oxygen	$+249 \text{ kJ mol}^{-1}$
First ionisation energy of Na	496 kJ mol^{-1}
First Electron affinity of Oxygen	-141 kJ mol^{-1}
First Electron affinity of Oxygen	798 kJ mol^{-1}
Enthalpy of formation of sodium oxide	-414 kJ mol^{-1}

QUESTION 3

Sum of the first and second electron affinities of O(g)

$$E_{\text{aff},1}^{\circ} + E_{\text{aff},2}^{\circ} = -141 + 798 = +657 \text{ kJ mol}^{-1}$$

Atomisation to form 1 mol O(g)

$$\Delta H_{\text{at}}^{\circ} = +249 \text{ kJ mol}^{-1}$$





Ionisation of 2 mol Na(g)

$$2 \times \Delta H_{\text{I1}}^{\circ} = +992 \text{ kJ mol}^{-1}$$

Atomisation to form 2 mol Na(g)

$$2 \times \Delta H_{\text{at}}^{\circ} = +214 \text{ kJ mol}^{-1}$$

Standard enthalpy change of formation of Na₂O(s)
 $\Delta H_f^{\circ}[\text{Na}_2\text{O(s)}] = ?$



Lattice enthalpy of sodium oxide,
 $\Delta H_{\text{lattice}}^{\circ}[\text{Na}_2\text{O(s)}] = ?$

QUESTION 4


Calculate the enthalpy change of formation of magnesium oxide

Enthalpy of atomisation of Mg	+147 kJ mol ⁻¹
Bond Enthalpy of Oxygen	+498 kJ mol ⁻¹
First ionisation energy of Mg	736 kJ mol ⁻¹
Second ionisation energy of Mg	1446 kJ mol ⁻¹
First electron affinity of Oxygen	-141 kJ mol ⁻¹
Second electron affinity of Oxygen	+791 kJ mol ⁻¹
Enthalpy of formation of MgO	-602 kJ mol ⁻¹

QUESTION 4



BORN HABER CYCLE FOR Al_2O_3



BORN-HABER CYCLES

If the cycles are examined it will be seen that:

the largest single contributions in each cycle are made by lattice energy and ionization energy and these two are always opposite in sign

the magnitude of the enthalpy of formation depends on the relative values of the lattice energy and the ionisation energy

if ionisation energy is too high to be supplied by lattice energy then such a compound will not form ionically

ENTHALPY CHANGES ON DISSOLVING


Why do ionic crystals dissolve in water, even though the ions in the lattice are strongly attracted to each other?

An ionic compound such as sodium chloride does not dissolve in non-polar solvents like hexane, but it will dissolve in a polar solvent like water. When one mole of sodium chloride dissolves in excess water to produce a very dilute solution under standard conditions, there is an enthalpy change of $+3.8 \text{ kJ mol}^{-1}$. This enthalpy change is described as the standard enthalpy change of solution, $\Delta H_{\text{solution}}$, of sodium chloride. The process can be summarised by the equation:



ENTHALPY CHANGES DURING DISSOLVING


When ionic compounds, like NaCl, dissolve in water the crystal lattice should be broken down and this requires input of energy. This energy is supplied through the energy of hydration.



ENTHALPY CHANGES DURING DISSOLVING

First of all, Na^+ and Cl^- ions must be removed from the solid NaCl crystals to form well-spaced ions in the gaseous state, $\text{Na}^+(\text{g})$ and $\text{Cl}^-(\text{g})$. This is the reverse of the lattice enthalpy (+787 kJ mol⁻¹).

In the second stage, gaseous $\text{Na}^+(\text{g})$ and $\text{Cl}^-(\text{g})$ ions are hydrated by polar water molecules forming a solution of sodium chloride, $\text{Na}^+(\text{aq})$ and $\text{Cl}^-(\text{aq})$. Under standard conditions, this process is the sum of the standard enthalpy changes of hydration of $\text{Na}^+(\text{g})$ and $\text{Cl}^-(\text{g})$.



ENTHALPY CHANGES DURING DISSOLVING


Enthalpy of Hydration:

The enthalpy change of hydration, ΔH_{hyd} , is the enthalpy change when 1 mole of a specified gaseous ion dissolves in sufficient water to form a very dilute solution.



Enthalpy of Solution:

The enthalpy change of solution, ΔH_{sol} , is the energy absorbed or released when 1 mole of an ionic solid dissolves in sufficient water to form a very dilute solution.



$$\Delta H_{\text{LE}} \text{NaCl} + \Delta H_{\text{soln}} \text{NaCl} = \Delta H_{\text{hyd}} \text{Na}^{+} + \Delta H_{\text{hyd}} \text{Cl}^{-}$$

$$\Delta H_{\text{LE}} + \Delta H_{\text{soln}} = \Delta H_{\text{hyd}}$$

$$\Delta H_{\text{soln}} = -\Delta H_{\text{LE}} + \Delta H_{\text{hyd}}$$


ENTHALPY OF SOLUTION

Draw a simple energy cycle & calculate the enthalpy of solution of sodium chloride from the following data

Lattice energy of sodium chloride = -776 kJ mol^{-1}


Hydration energy of $\text{Na}^+(\text{g})$ = -390 kJ mol^{-1}

Hydration energy of $\text{Cl}^-(\text{g})$ = -381 kJ mol^{-1}



QUESTION 1

- (a) What is meant by the term enthalpy change of hydration, ΔH_{hyd} ?
- (b) Write an equation that represents the of the ΔH_{hyd} Mg^{2+} ion.
- (c) Suggest a reason why ΔH_{hyd} of the Mg^{2+} ion is greater than of the Ca^{2+} ion.
- (d) Suggest why it is impossible to determine the enthalpy change of hydration of the oxide ion, O^{2-} .




If the energy of hydration of ions outweighs the lattice energy the compound dissolves exothermally. If the energy of hydration is much lower than the lattice energy of the compound, the compound will not dissolve. If the energy of hydration is slightly less than the lattice energy, the little extra energy required for the breakdown of the lattice is obtained from the thermal energy of the solvent. Hence the process would be endothermic resulting in a drop in temperature.

Most salts dissolve endothermically. Hence an increase of temperature favors the equilibrium point to shift forwards resulting in an increase in the solubility.

ENTHALPY OF SOLUTION

By constructing relevant thermochemical cycles, use the above data to calculate a value for $\Delta H_{\text{sol}}(\text{MgCl}_2(\text{s}))$



QUESTION 2

Iron(III) chloride redissolves




Use the following data to calculate the standard enthalpy change for this process.

species	$\Delta H_f^\circ / \text{kJ mol}^{-1}$
$\text{FeCl}_3(\text{s})$	−399.5
$\text{Fe}^{3+}(\text{aq})$	−48.5
$\text{Cl}^-(\text{aq})$	−167.2

QUESTION 2

By constructing relevant thermochemical cycles, use the above data to calculate a value for $\Delta H_{\text{sol}}(\text{MgCl}_2(\text{s}))$



$$(-48.5) + 3(-167.2) = -399.5 + x$$

$$-150.1$$